

Definable obstruction theory

Tomohiro Kawakami¹ and Ikumitsu Nagasaki²

¹Department of Mathematics, Faculty of Education, Wakayama University,
Sakaedani Wakayama 640-8510, Japan

²Department of Mathematics, Kyoto Prefectural University of Medicine,
13 Nishi-Takatsukaso-Cho, Taishogun Kita-ku, Kyoto 603-8334, Japan

Received September 30, 2014

Abstract

Let $\mathcal{N} = (R, +, \cdot, <, \dots)$ be an o-minimal expansion of the standard structure of a real closed field R . In this paper, we consider an obstruction theory in the definable category of \mathcal{N} .

2010 *Mathematics Subject Classification.* 55N20, 03C64.

Keywords and Phrases. Obstruction theory, o-minimal, real closed fields.

1 . Introduction.

Obstruction theory addresses several types of problems(see chap. 7 [2]). Let (X, A) be a *CW* pair and Y a topological space. One of these problems is Extension Problem.

Problem 1.1. *Suppose that $f : A \rightarrow Y$ is a continuous map. When does f extend to all of X ?*

Let $\mathcal{N} = (R, +, \cdot, <, \dots)$ be an o-minimal expansion of the standard structure of a real closed field R . General references on o-minimal structures are [3], [5], see also [9]. Examples and constructions of them can be seen in [4], [6], [7].

In this paper, we consider an obstruction theory in the definable category of \mathcal{N} . Everything is considered in \mathcal{N} , a definable map is assumed to be continuous and $I = \{x \in R \mid 0 \leq x \leq 1\}$.

Theorem 1.2. *Let (X, A) be a relative definable CW complex, $n \geq 1$, and Y a de-*

finably connected n -simple definable set. Let $g : X_n \rightarrow Y$ be a definable map.

- (1) *There exists a cellular cocycle $\theta(g) \in C^{n+1}(X, A, \pi_n(Y))$ which vanishes if and only if g extend to a definable map $X_{n+1} \rightarrow Y$.*
- (2) *The cohomology class $[\theta(g)] \in H^{n+1}(X, A, \pi_n(Y))$ vanishes if and only if the restriction $g|_{X_{n-1}} : X_{n-1} \rightarrow Y$ extend to a definable map $X_{n+1} \rightarrow Y$.*

2 . Preliminaries.

Let $X \subset R^n$ and $Y \subset R^m$ be definable sets. A continuous map $f : X \rightarrow Y$ is *definable* if the graph of f ($\subset X \times Y \subset R^n \times R^m$) is a definable set. A definable map $f : X \rightarrow Y$ is a *definable homeomorphism* if there exists a definable map $h : Y \rightarrow X$ such that $f \circ h = id_Y, h \circ f = id_X$. A definable subset X of R^n is *definably compact* if for every definable map $f : (a, b)_R \rightarrow X$, there

exist the limits $\lim_{x \rightarrow a+0} f(x)$, $\lim_{x \rightarrow b-0} f(x)$ in X , where $(a, b)_R = \{x \in R \mid a < x < b\}$, $-\infty \leq a < b \leq \infty$. A definable subset X of R^n is definably compact if and only if X is closed and bounded ([8]). Note that if X is a definably compact definable set and $f : X \rightarrow Y$ is a definable map, then $f(X)$ is definably compact.

If R is the field \mathbb{R} of real numbers, then for any definable subset X of \mathbb{R}^n , X is compact if and only if it is definably compact. In general, a definably compact definable set is not necessarily compact. For example, if $R = \mathbb{R}_{alg}$, then $[0, 1]_{\mathbb{R}_{alg}} = \{x \in \mathbb{R}_{alg} \mid 0 \leq x \leq 1\}$ is definably compact but not compact.

Recall existence of definable quotient and properties of dimensions of definable sets.

Theorem 2.1. (*Existence of definable quotient (e.g. 10. 2.14 [3])*). If X is a definable set and A is a definably compact definable subset of X , then the set obtained by collapsing A to a point exists a definable set.

Proposition 2.2 (e.g. 4.1.3 [3]). (1) If $X \subset Y \subset R^n$, then $\dim X \leq \dim Y \leq n$.
(2) If $X \subset R^n$, $Y \subset R^m$ are definable sets and there is a definable bijection between X and Y , then $\dim X = \dim Y$.

Let $(X, A), (Y, B)$ be two pairs of definable sets. Two definable maps $f, h : (X, A) \rightarrow (Y, B)$ is *definably homotopic relative to A* if there exists a definable map $H : (X \times I, A \times I) \rightarrow (Y, B)$ such that $H(x, 0) = f(x)$, $H(x, 1) = g(x)$ for all $x \in X$ and $H(x, t) = f(x)$, $(x, t) \in A \times I$. The *α -minimal homotopy set* $[(X, A), (Y, B)]$ of (X, A) and (Y, B) is the set of homotopy classes of definable maps from (X, A) to (Y, B) . If $A = \emptyset, B = \emptyset$, then we simply write $[X, Y]$ instead of $[(X, A), (Y, B)]$.

Let $D^n = \{(x_1, \dots, x_n) \in R^n \mid x_1^2 + \dots + x_n^2 \leq 1\}$, $S^{n-1} = \{(x_1, \dots, x_n) \in R^n \mid x_1^2 + \dots + x_n^2 = 1\}$. Then D^n is the closed unit disk of R^n and S^{n-1} is the unit sphere of R^n .

We now define relative CW complexes in the definable category. To reserve definability, we consider the case where finitely many cells attached.

Definition 2.3. Let X be a definable set and A a definable closed subset of X . We say that X is obtained from A by attaching n -cells $\{e_i^n\}_{i=1}^{k_n}$ if the following four conditions satisfy.

(1) For each i , e_i^n is a definable subset of X , called an n -cell.

(2) $X = A \cup \bigcup_{i=1}^{k_n} e_i^n$.

(3) Letting ∂e_i^n denote the intersection of e_i^n and A , $e_i^n - \partial e_i^n$ is disjoint from $e_j^n - \partial e_j^n$ for $i \neq j$.

(4) For each i , there exists a surjective definable map $\phi_i^n : (D^n, S^{n-1}) \rightarrow (e_i^n, \partial e_i^n)$, called the *characteristic map* of e_i^n , such that the restriction of ϕ_i of the interior $\text{Int } D^n$ of D^n is a definable homeomorphism onto $e_i^n - \partial e_i^n$. The restriction of the characteristic map of S^{n-1} is the *attaching map* of e_i^n .

Definition 2.4. A *relative definable CW complex* (X, A) is a definable set X , a definable closed set A and a sequence of definable closed subset X_n , $n = -1, 0, 1, 2, \dots$ called the *relative n -skeleton* such that

(1) $X_{-1} = A$ and X_n is obtained from X_{n-1} by attaching n -cells.

(2) $X = \bigcup_{i=-1}^{\dim X} X_i$.

The smallest n such that $X = X_n$ is called the *dimension* $\dim(X, A)$ of (X, A) . If A is a definable CW complex, we say that (X, A) is a *definable CW pair*. If $A = \emptyset$, then X is called a *definable CW complex*, and X_n is called the *n -skeleton* of X .

Remark that in Definition 2.4, the maximum dimension of attaching cells to A does not exceed $\dim X$ and $\dim A \leq \dim X$ because Proposition 2.2.

Let Y be a definable set and $y_0 \in Y$. The *α -minimal homotopy group of dimension n* , $n \geq 1$ (see [1]) is the set $\pi_n(Y, y_0) = [(I^n, \partial I^n), (Y, y_0)] = [(S^n, x_0), (Y, y_0)]$, where ∂I^n denote the boundary of I^n and $x_0 = (0, \dots, 0, 1)$. We define $\pi_0(Y, y_0)$ as the set of definably connected components of Y .

A definable set Y is *definably arcwise connected* if for every two points $x, y \in Y$, there exists a definable map $f : I \rightarrow Y$ such that $x = f(0)$ and $y = f(1)$. Note that Y is

definably connected if and only if it is definably arcwise connected. In this case, for any $y_0, y_1 \in Y$ and $n \geq 1$, $\pi_n(Y, y_0)$ is isomorphic to $\pi_n(Y, y_1)$ and we denote it $\pi_n(Y)$.

For $n \geq 1$, a definably connected definable set is *definably n -connected* if $\pi_i(Y) = 0$ for each $1 \leq i \leq n$.

Lemma 2.5. *Let Y be a definably connected definable set. If $\pi_{n-1}(Y) = 0$, then for every definable map $h : S^{n-1} \rightarrow Y$, there exists a definable map $H : D^n \rightarrow Y$ with $H|_{S^{n-1}} = h$.*

Proof. For $i \geq 1$, since Y is definably connected, $\pi_i(Y) \rightarrow [S^i, Y], [h] \rightarrow [h]$ is bijective. Thus h is definably homotopic to a constant map $C : S^{n-1} \rightarrow Y, C(x) = c$. Hence there exists a definable map $\phi : S^{n-1} \times I \rightarrow Y$ such that $\phi(x, 0) = c, \phi(x, 1) = h(x)$ for all $x \in S^{n-1}$. Collapsing $S^{n-1} \times \{0\}$ to a point, by Theorem 2.1, we have the cone CS^{n-1} which is definably homeomorphic to D^n and a definable map $H : D^n \rightarrow Y$ with $H|_{S^{n-1}} = h$. \square

Proposition 2.6. *If Y is definably $(n-1)$ -connected, $f : A \rightarrow Y$ is a definable map, $\dim(X, A) \leq n$ and $n \geq 1$, then there exists a definable map $F : X \rightarrow Y$ with $F|_A = f$.*

Proof. If $i = 0$, then we may assume that $X_0 = A \cup e_1^0 \cup \dots \cup e_{r_0}^0$, $e_1^0, \dots, e_{r_0}^0$ denote the 0-cells of (X, A) . For each e_j^0 , defining the image of e_j^0 , there exists a definable map $f_0 : X_0 \rightarrow Y$ extending f .

We may assume that $X_i = X_{i-1} \cup e_1^i \cup \dots \cup e_{r_i}^i$, $e_1^i, \dots, e_{r_i}^i$ denote the i -cells of (X, A) . By assumption, there exists a definable map $h_j : \partial e_j^i \rightarrow Y$. Since ∂e_j^i is definably homeomorphic to S^{n-1} and by Lemma 2.5, we have a definable map $H_j : e_j^i \rightarrow Y$ with $H_j|_{\partial e_j^i} = h_j$. Using H_j , we obtain a definable map F with $F|_A = f$. \square

Let X be a definably connected definable set and $n \geq 1$. As in the topological setting, $\pi_1(X)$ acts on $\pi_n(X)$. We say that a definably connected definable set X is *n -simple* if the $\pi_1(X)$ action on $\pi_n(X)$ is trivial. Since the $\pi_1(X)$ action on $\pi_1(X)$ is

$\pi_1(X) \times \pi_1(X) \rightarrow \pi_1(X), (h_1, h_2) \mapsto h_1 h_2 h_1^{-1}$, X is 1-simple if and only if $\pi_1(X)$ is abelian.

Let X be a definable CW complex, A a definable subcomplex of X , $n \geq 1$ and Y a definably connected n -simple definable set. We define the cohomology group $H^n(X, A, \pi_n(Y))$ as follows. Remark that $[S^n, Y] = \pi_n(Y)$ because Y is n -simple.

We define the n -dimensional chain complex $C_n(X, A)$ to be $H_n(X_n, X_{n-1})$. Let $i_{n-1} : X_{n-1} \rightarrow X_n, j_n : (X_n, \emptyset) \rightarrow (X_n, X_{n-1})$ be inclusions. As in the topological setting, we have an exact sequence

$$\dots \rightarrow H_n(X_n, X_{n-1}) \xrightarrow{\partial'_n} H_{n-1}(X_{n-1}) \xrightarrow{i_{n-1}^*} \dots$$

$$H_{n-1}(X_n) \xrightarrow{j_n^*} H_{n-1}(X_n, X_{n-1}) \rightarrow \dots$$

The boundary operator $\partial_n : H_n(X_n, X_{n-1}) \rightarrow H_{n-1}(X_{n-1}, X_{n-2})$ is $j_{n-1}^* \circ \partial'_n$. We define the n -dimensional cochain complex $C^n(X, A) = \text{Hom}_{\mathbb{Z}}(C_n(X, A), \pi_n(Y))$ and the coboundary operator $\delta_n : C^n(X, A) \rightarrow C^{n+1}(X, A), (\delta f)c = f(\partial c)$.

Let (X, A) be a relative definable CW complex, $n \geq 1$, and Y a definably connected n -simple definable set. Let $g : X_n \rightarrow Y$ be a definable map.

Let e_i^{n+1} be an $(n+1)$ -cell and $\phi_i : (D^{n+1}, S^n) \rightarrow (e_i^{n+1}, \partial e_i^{n+1}) \subset (X_{n+1}, X_n)$ the characteristic map of e_i^{n+1} . Composing $f_i = \phi_i|_{S^n}$ with $g : X_n \rightarrow Y$, we have an element $[g \circ f_i] \in [S^n, Y] = \pi_n(Y)$. We define the obstruction cochain $\theta^{n+1}(g) \in C^{n+1}(X, A, \pi_n(Y))$ on the basis of $(n+1)$ -cells by the formula $\theta^{n+1}(g)(e_i^{n+1}) = [g \circ f_i]$ and extend by linearity.

In the rest of this section, we prove the o-minimal cellular approximation theorem

Theorem 2.7 (O-minimal cellular approximation theorem). *Let $(X, A), (Y, B)$ be definable CW pairs and $f : (X, A) \rightarrow (Y, B)$ a definable map. Then there exists a definable map $g : (X, A) \rightarrow (Y, B)$ such that f is definably homotopic to g relative to A and for any nonnegative integer n , $g(X'_n) \subset Y'_n$, where X'_n (resp. Y'_n) denotes the union of the n -skeleton X_n (resp. Y_n) of X (resp. Y) and A (resp. B).*

Lemma 2.8 (O-minimal homotopy extension lemma [1]). *Let X, Z, A be definable sets with $A \subset X$ closed in X . Let $f : X \rightarrow Z$ be a definable map and $H : A \times I \rightarrow Z$ a definable homotopy such that $H(x, 0) = f(x), x \in A$. Then there exists a definable homotopy $F : X \times I \rightarrow Z$ such that $F(x, 0) = f(x), x \in X$ and $F|A \times I = H$.*

By the above lemma, we have the following o-minimal homotopy extension theorem.

Theorem 2.9. *Let (X, A) be a definable CW pair. Let $f : X \rightarrow Y$ be a definable map and $H : A \times I \rightarrow Y$ a definable homotopy with $H(x, 0) = f(x), x \in A$. Then there exists a definable homotopy $F : X \times I \rightarrow Y$ such that $F(x, 0) = f(x), x \in X$ and $F|A \times I = H$.*

To prove Theorem 2.7, we prepare three claims.

Claim 2.10. *Let (Z, C) be a definable CW pair. For any definable map $g : D^q \rightarrow Z$ with $g(S^{q-1}) \subset \overline{Z^{q-1}}$, there exists a definable map $g' : D^q \rightarrow Z$ such that $g \simeq g' \text{ rel } S^{q-1}$ and $g'(D^q) \subset \overline{Z^q}$, where $\overline{Z^{q-1}} = Z^q \cup C$.*

Proof. Let n be the maximum dimension of cells not contained in C . We may assume that $n > q$ and proceed by induction on the number of such n -cells. Let $\phi : (D^n, S^{n-1}) \rightarrow (Z, \overline{Z^{n-1}})$ be the characteristic map of an n -cell e . Let $D_1^n, (D_2^n)$ be the closed ball of center 0 with radius $\frac{1}{3}, (\frac{2}{3})$, respectively. Put $U = \phi(D^n - D_1^n) \cup (Z - e), V = \phi(\text{Int } D_2^n), z_0 = \phi(0)$, where $\text{Int } D_2^n$ denotes the interior of D_2^n . Then $U \cup V = Z$. Taking a refinement of D^q , every simplex $|s|$ of it is contained in $g^{-1}(U)$ or $g^{-1}(V)$. Let $E_1 = \cup_{|s| \cap g^{-1}(z_0) \neq \emptyset} |s|, E_2 = \cup_{|s| \cap g^{-1}(z_0) = \emptyset} |s|$. Then $g(E_1) \subset V, g(E_1 \cap E_2) \subset V - \{z_0\}$. Thus we have a definable map $\phi^{-1} \circ g : E_1 \cap E_2 \rightarrow \text{Int } D_2^n - \{0\}$. Since $\text{Int } D_2^n - \{0\}$ is definably homotopy equivalent to S^{n-1} and S^{n-1} is $(n-2)$ -connected, there exists a definable map $h : E_1 \rightarrow \text{Int } D_2^n - \{0\}$ with $h|E_1 \cap E_2 = \phi^{-1} \circ g$. Define a definable homotopy $h_t : E_1 \rightarrow \text{Int } D_2^n$ by $h_t(x) =$

$(1-t)\phi^{-1} \circ g(x) + th(x)$. Then h_t is a definable homotopy between $\phi^{-1} \circ g$ and h relative to $E_1 \cap E_2$. Define a definable homotopy $h'_t : D^q \rightarrow Z$ by $h'_t|E_1 = \phi^{-1} \circ g, h'_t|E_2 = g|E_2$. Then h'_t is a definable homotopy between g and h'_1 relative to S^{q-1} and $h'_1(D^q) \subset Z - \{z_0\}$. Taking a definable retraction $r : Z - \{z_0\} \rightarrow Z - e, h'_1 \simeq r \circ h'_1 \text{ rel } S^{q-1} : D^q \rightarrow Z - \{z_0\}$. Let $g'' = r \circ h'_1$. Then $g \simeq g'' \text{ rel } S^{q-1} : D^q \rightarrow Z, g''(D^q) \subset Z - e$. By the inductive hypothesis, there exists a definable map g' such that $g'' \simeq g' \text{ rel } S^{q-1} : D^q \rightarrow Z - e, g'(D^q) \subset \overline{Z^q}$. \square

Claim 2.11. *For any definable map $f : (\overline{X^q}, \overline{X^{q-1}}) \rightarrow (Y, \overline{Y^{q-1}})$, there exists a definable map $g : (\overline{X^q}, \overline{X^{q-1}}) \rightarrow (Y, \overline{Y^{q-1}})$ such that $f \simeq g \text{ rel } \overline{X^{q-1}}$ and $g(\overline{X^q}) \subset \overline{Y^q}$.*

Proof. Let e be a q -cell not contained in A . Since $f(\overline{e})$ is definably compact, there exists a finite subcomplex Z of Y with $f(\overline{e}) \subset Z$. Put $C = Z \cap \overline{Y^{q-1}}$. Then $f(e^r) \subset C$, where e^r denotes the boundary of e . Let $\phi : (D^q, S^{q-1}) \rightarrow (\overline{e}, e^r)$ be the characteristic map of e . Applying Claim 2.10 to $f \circ \phi : (D^q, S^{q-1}) \rightarrow (Z, C)$, there exists a definable map g' such that $f \circ \phi \simeq g' \text{ rel } S^{q-1}, g'(D^q) \subset \overline{Z^q}$. Then $g = g' \circ \phi$ is the required map. \square

Claim 2.12. *For any definable map $f : (X, A) \rightarrow (Y, B)$, there exists a definable homotopy $H_q : (X, A) \times [0, 1]_R \rightarrow (Y, B)$ such that:*

- (1) $H_0(x, t) = f(x)$ for all $x \in X$.
- (2) $H_q(x, 0) = H_{q+1}(x, 0)$ for all $x \in X$.
- (3) $H_q(x, t) = (x, t)$ for all $(x, t) \in \overline{X^q} \times [0, 1]_R$.
- (4) $H_q(\overline{X^q} \times \{1\}) \subset \overline{Y}$.

Proof. Let $H_0(x, t) = f(x)$ for $(x, t) \in X \times [0, 1]_R$. Assume we have H_{q-1} . Since $H_{q-1}(\overline{X^{q-1}} \times \{1\}) \subset \overline{Y^{q-1}}$ and by Claim 2.11, there exists a definable homotopy $H'_q \text{ rel } \overline{X^{q-1}} : (\overline{X^q}, \overline{X^{q-1}}) \times [0, 1]_R \rightarrow (\overline{Y^q}, \overline{Y^{q-1}})$ such that $H'_q|_{\overline{X^q} \times \{0\}} = H_{q-1}|_{\overline{X^q} \times \{1\}}, H'_q(\overline{X^q} \times \{1\}) \subset \overline{Y^q}$. By Lemma 2.8, there exists a definable homotopy $H_q : X \times [0, 1]_R \rightarrow Y$

such that $H_q|X \times \{0\} = H_{q-1}|X \times \{1\}$, $H_q|\overline{X}^q \times [0, 1]_R = H'_q$, and H_q satisfies (1)-(4). \square

Proof of Theorem 2.7. Let $q = \dim X$. By Claim 2.12, we have a definable homotopy H_q . Then the definable map $g : (X, A) \rightarrow (Y, B)$ defined by $g(x) = H_q(x, 1)$ is the required map. \square

3. Proof of Theorem 1.2.

Lemma 3.1. *Let i be the inclusion $X_n \rightarrow X_{n+1}$ and $x_0 \in X_n$. Then $i_* : \pi_1(X_n, x_0) \rightarrow \pi_1(X_{n+1}, x_0)$ is surjective if $n = 1$ and an isomorphism $n > 1$.*

Proof. Let $n \geq 1$ and $\alpha : S^1 \rightarrow X_{n+1}$ a definable map. By Theorem 2.7, there exists a definable map $\alpha' : S^1 \rightarrow X_1 \subset X_n$ such that α is definably homotopic to α' . Since $i_*([\alpha']) = [\alpha]$, i_* is surjective.

Assume $n \geq 2$ and $i_*([\alpha]) = 0$. Then $\alpha : S^1 \rightarrow X_{n+1}$ is null homotopic and there exists a definable map $H : S^1 \times [0, 1]_R \rightarrow X_{n+1}$ such that $H(-, 0) = \alpha$, $H(-, 1) = c$, where c denotes a constant map. By Theorem 2.7 and since $S^1 \times [0, 1]_R$ is a 2-dimensional definable set, there exists a definable map $H' : S^1 \times [0, 1]_R \rightarrow X_2$ such that H is definably homotopic to H' relative to $S^1 \times \{0, 1\}$. Thus $[\alpha] = 0$ and i_* is injective. \square

Lemma 3.2. *If $k \leq n, n > 1$ and $x_0 \in X_n$, then $\pi_k(X_{n+1}, X_n, x_0) = 0$.*

Proof. Consider an exact sequence $\cdots \rightarrow \pi_k(X_n, x_0) \rightarrow \pi_k(X_{n+1}, x_0) \rightarrow \pi_k(X_{n+1}, X_n, x_0) \rightarrow \pi_{k-1}(X_n, x_0) \rightarrow \pi_{k-1}(X_{n+1}, x_0) \rightarrow \cdots$. We prove that $i_{*k} : \pi_k(X_n, x_0) \rightarrow \pi_k(X_{n+1}, x_0)$ is surjective and $i_{*k-1} : \pi_{k-1}(X_n, x_0) \rightarrow \pi_{k-1}(X_{n+1}, x_0)$ is injective.

Let $\alpha : S^k \rightarrow X_{n+1}$ be a definable map. Then by Theorem 2.7, there exists a definable map $\alpha' : S^k \rightarrow X_k$ such that α is definably homotopic to α' . Then $i_{*k} : \pi_k(X_n, x_0) \rightarrow \pi_k(X_{n+1}, x_0)$ is surjective.

Assume $i_{*k-1}([\alpha]) = 0$. Then $\alpha : S^{k-1} \rightarrow X_{n+1}$ is null homotopic and there exists a definable map $H : S^{k-1} \times [0, 1]_R \rightarrow X_{n+1}$ such that $H(-, 0) = \alpha$, $H(-, 1) = c$. By

Theorem 2.7 and since $S^{k-1} \times [0, 1]$ is a k -dimensional definable set, there exists a definable map $H' : S^{k-1} \times [0, 1]_R \rightarrow X_k \subset X_n$ such that H is definably homotopic to H' relative to $S^{k-1} \times \{0, 1\}$. Thus $[\alpha] = 0$ and i_{*k-1} is injective.

By the above results and exactness, we have the lemma. \square

The following is the o-minimal relative Hurewicz theorem.

Theorem 3.3 (5.4 [1]). *Let (X, A, x_0) be a definable pointed pair and $n \geq 2$. Suppose that $\pi_r(X, A, x_0) = 0$ for any $1 \leq r \leq n - 1$. Then the o-minimal Hurewicz homomorphism $h_n : \pi_n(X, A, x_0) \rightarrow H_n(X, A)$ is surjective and its kernel is the subgroup generated by $\{\beta_{[u]}([f])[f]^{-1}[u] \in \pi_1(A, x_0), [f] \in \pi_n(X, A, x_0)\}$. In particular, h_n is an isomorphism for $n \geq 3$.*

Put $\pi_{n+1}^+(X_{n+1}, X_n) = \pi_{n+1}(X_{n+1}, X_n) / \ker h_n$. Let $g : X_n \rightarrow Y$ be a definable map and $\pi : \pi_{n+1}(X_{n+1}, X_n) \rightarrow \pi_{n+1}^+(X_{n+1}, X_n)$ denote the projection.

Lemma 3.4. *There exists a factorization $\overline{g_* \circ \partial} : \pi_{n+1}^+(X_{n+1}, X_n) \rightarrow \pi_n(Y)$ such that $\pi \circ \overline{g_* \circ \partial} = g_* \circ \partial$.*

Proof. If $\alpha \in \pi_1(X_n)$, then $\partial(\alpha x) = a\partial x$. Since Y is n -simple, for any $z \in \pi_n(X_n)$, $g_*(\alpha z) = g_*(\alpha)g_*(z) = g_*(z)$. \square

By Lemma 3.4, we can define the composition map $C_{n+1}(X, A) = H_{n+1}(X_{n+1}, X_n) \xrightarrow{h^{-1}}$

$\pi_{n+1}^+(X_{n+1}, X_n) \xrightarrow{g_* \circ \partial} \beta \pi_n(Y)$, where $h : \pi_{n+1}^+(X_{n+1}, X_n) \rightarrow H_{n+1}(X_{n+1}, X_n)$ denotes the Hurewicz isomorphism. This composition map defines another cochain in $Hom_{\mathbb{Z}}(C_{n+1}(X, A), \pi_n(Y))$ which we again denote by $\theta^{n+1}(g)$.

Proposition 3.5. *The two definitions of $\theta^{n+1}(g)$ coincide.*

Proof. For an $(n + 1)$ -cell e_i^{n+1} , let $\phi_i : (D^{n+1}, S^n) \rightarrow (X_{n+1}, X_n)$ be the characteristic map of e_i^{n+1} . We define a map $(\phi_i \vee u) \circ q : (D^{n+1}, S^n, p) \rightarrow (X_{n+1}, X_n, x_0)$ as the

composition of a map $q : (D^{n+1}, S^n, p) \rightarrow (D^{n+1} \vee I, S^n \vee I, p)$ and a map $D^{n+1} \vee I \xrightarrow{\phi_i \vee u} X_{n+1}$, where u is a definable path in X_n to the base point x_0 . Then $(\phi_i \vee u) \circ q$ is definably homotopic to the characteristic map ϕ_i . Hence $h((\phi_i \vee u) \circ q)$ is the generator of $H_{n+1}(X_{n+1}, X_n)$ represented by the cell e_i^{n+1} and $(\phi_i \vee u) \circ q$ represents the element $h^{-1}(e_i^{n+1})$ in $\pi_{n+1}^+(X_{n+1}, X_n)$. By definition, $\partial((\phi_i \vee u) \circ q) \in \pi_n(X_n)$ is represented by the composition of the map $\bar{q} : S^n \rightarrow S^n \vee I$ obtained by restricting the map q to the boundary and the attaching map $f_i = \phi_i|_{S^n}$ together with a definable path u to x_0 : $\partial((\phi_i \vee u) \circ q) = (f_i \vee u) \circ \bar{q} : S^n \rightarrow X_n$. By the second definition, $\theta(g)(e_i^{n+1}) = g \circ (f_i \vee u) \circ \bar{q} = (g_i \circ f_i \vee g \circ u) \circ \bar{q}$. Moreover this is equal to $[f_i] \in [S^n, Y] = \pi_n(Y)$, which is the first definition of $\theta(g)(e_i^{n+1})$. \square

Theorem 3.6. *The obstruction cochain $\theta^{n+1}(g)$ is a cocycle.*

Proof. Consider the following commutative diagram.

$$\begin{array}{ccc}
 \pi_{n+2}(X_{n+2}, X_{n+1}) & \rightarrow & H_{n+2}(X_{n+2}, X_{n+1}) \\
 \downarrow & & \downarrow \\
 \pi_{n+1}(X_{n+1}) & \rightarrow & H_{n+1}(X_{n+1}) \\
 \downarrow & & \downarrow \\
 \pi_{n+1}(X_{n+1}, X_n) & \rightarrow & H_{n+1}(X_{n+1}, X_n) \\
 \downarrow & & \downarrow \theta(g) \\
 \pi_n(X_n) & \xrightarrow{g_*} & Hn(Y_n)
 \end{array}$$

The unlabeled horizontal arrows are the Hurewicz maps and the unlabeled vertical arrows are obtained from homotopy or homology exact sequences of the pair (X_{n+2}, X_{n+1}) and (X_{n+1}, X_n) .

The composition of the bottom two vertical maps on the left are zero because they occur in the homotopy exact sequence of the pair (X_{n+1}, X_n) . Since $\delta\theta(g)$ is the composition of all the right vertical maps, $\delta\theta^{n+1}(g)(x) = \theta^{n+1}(g)(\partial x) = 0$. Thus $\theta^{n+1}(g)$ is a cocycle. \square

By a way to similar to the topological category, we have the following proposition.

Proposition 3.7. *If X is a definable CW complex, then $X \times I$ is a definable CW complex.*

Theorem 3.8. *Let (X, A) be a relative definable CW complex, Y a definably connected n -simple definable set and $g : X_n \rightarrow Y$ a definable map.*

(1) $\theta^{n+1}(g) = 0$ if and only if there exists a definable map $\tilde{g} : X_{n+1} \rightarrow Y$ extending g .

(2) $[\theta^{n+1}(g)] = 0$ if and only if there exists a definable map $\tilde{g} : X_{n+1} \rightarrow Y$ extending $g|_{X_{n-1}}$.

Lemma 3.9. *Let $f_0, f_1 : X_n \rightarrow Y$ be definable maps such that $f_0|_{X_{n-1}}$ is definably homotopic to $f_1|_{X_{n-1}}$. Then there exists a difference cochain $d \in C^n(X, A, \pi_n(Y))$ such that $\delta d = \theta^{n+1}(f_0) - \theta^{n+1}(f_1)$.*

Proof. Let $\hat{X} = X \times I$, $\hat{A} = A \times I$. Then (\hat{X}, \hat{A}) is a relative definable CW complex with $\hat{X}^k = X_k \times \partial I \cup X_{k-1} \times I$. Take a definable homotopy H between f_0 and f_1 . Hence a definable map $\hat{X}_n \rightarrow Y$ is obtained from $f_0, f_1 : X_n \rightarrow Y$ and a definable homotopy $G = H|_{X_{n-1} \times I} : X_{n-1} \times I \rightarrow Y$. Thus we have the cocycle $\theta(f_0, G, f_1) \in C^{n+1}(\hat{X}, \hat{A}, \pi_n(Y))$ which obstructs finding an extension of $f_0 \cup G \cup f_1$ to \hat{X}_{n+1} . we define the difference cochain $d(f_0, G, f_1) \in C^n(X, A, \pi_n(Y))$ by restricting to cells of the form $e^n \times I$, that is $d(f_0, G, f_1)(e_i^n) = (-1)^{n+1} \theta(f_0, G, f_1)(e_i^n \times I)$ for each n -cell e_i^n of X . Since $\theta(f_0, G, f_1)$ is a cocycle, $0 = (\delta\theta(f_0, G, f_1))(e_i^{n+1} \times I) = \theta(f_0, G, f_1)(\partial((e_i^{n+1} \times I))) = \theta(f_0, G, f_1)(\partial(e_i^{n+1} \times I) + (-1)^{n+1}(\theta(f_0, G, f_1)(e_i^{n+1} \times \{1\}) - \theta(f_0, G, f_1)(e_i^{n+1} \times \{0\}))) = (-1)^{n+1}(\delta(d(f_0, G, f_1))(e_i^{n+1}) + \theta(f_1)(e_i^{n+1}) - \theta(f_0)(e_i^{n+1}))$. Thus $\delta d = \theta^{n+1}(f_0) - \theta^{n+1}(f_1)$. \square

Proposition 3.10. *Let $f_0 : X_n \rightarrow Y$ be a definable map, $G : X_{n-1} \times I \rightarrow Y$ a definable homotopy such that $G(-, 0) = f_0|_{X_{n-1}}$ and $d \in C^n(X, A, \pi_n(Y))$. Then there exists a definable map $f_1 : X_n \rightarrow Y$ such that $G(-, 1) = f_1|_{X_{n-1}}$ and $d = d(f_0, G, f_1)$.*

To prove Proposition 3.10, we need the following lemma.

Lemma 3.11. *For any definable map $f : D^n \times \{0\} \cup S^{n-1} \times I \rightarrow Y$ and for any definable homotopy class $\alpha \in [\partial(D^n \times I), Y]$, there exists a definable map $F : \partial(D^n \times I) \rightarrow Y$ such that $F|_{D^n \times \{0\} \cup S^{n-1} \times I} = f$ and $[F] = [\alpha]$.*

Proof. Take a definable map $K : \partial(D^n \times I) \rightarrow Y$ with $[K] = [\alpha]$. Let $D = D^n \times \{0\} \cup S^{n-1} \times I$. Then D is definably contractible and $K|_D$ and f are null homotopic. Thus $K|_D$ and f are definably homotopic. Applying Theorem 2.7 to $(\partial(D^n \times I), D)$, there exists an extension $H : \partial(D^n \times I) \times I \rightarrow Y$. Hence $F = H(-, 1)$ is the required definable map. \square

Proof of Proposition 3.10. Let e_i^n be an n -cell of X_n and $\phi : (D^n, S^{n-1}) \rightarrow (X_n, X_{n-1})$ the characteristic map of e_i^n . Applying Lemma 3.11 to $f = f_0 \circ \phi_i \cup G \circ (\phi_i|_{S^{n-1}} \times id_I)$ and $\alpha = d(e_i^n)$, we have a definable map F_i . We define $f_1 : X_n \rightarrow Y$ on the n -cells by $f_1(\phi_i(x)) = F_i(x, 1)$. Then $d(f_0, G, f_1)(e_i^n) = d(e_i^n)$. \square

Proof of Theorem 3.8. We now prove that if $g : X_n \rightarrow Y$ and $\theta(g)$ is a coboundary δd , then $g|_{X_{n-1}}$ extends to X_n . Applying Proposition 3.10 to g , d and the stationary homotopy $((x, t) \mapsto g(x))$ from $g|_{X_{n-1}}$ to itself, there exists a definable map $g' : X_n \rightarrow Y$ such that $g'|_{X_{n-1}} = g|_{X_{n-1}}$ and $\delta d = \theta(g) - \theta(g')$. Since $\theta(g') = 0$, g' extends to X_{n+1} . \square

References

- [1] E. Baro and M. Otero, *On o-minimal homotopy groups*, Q. J. Math. **61** (2010), 275–289.
- [2] J.F. Davis and P. Kirk, *Lecture notes in algebraic topology*, Graduate Studies in Mathematics, **35**, American Mathematical Society, Providence, RI, (2001).
- [3] L. van den Dries, *Tame topology and o-minimal structures*, Lecture notes series **248**, London Math. Soc. Cambridge Univ. Press (1998).
- [4] L. van den Dries, A. Macintyre and D. Marker, *The elementary theory of restricted analytic field with exponentiation*, Ann. of Math. **140** (1994), 183–205.
- [5] L. van den Dries and C. Miller, *Geometric categories and o-minimal structures*, Duke Math. J. **84** (1996), 497–540.
- [6] L. van den Dries and P. Speissegger, *The real field with convergent generalized power series*, Trans. Amer. Math. Soc. **350** (1998), 4377–4421.
- [7] C. Miller, *Expansion of the field with power functions*, Ann. Pure Appl. Logic **68**, (1994), 79–94.
- [8] Y. Peterzil and C. Steinhorn, *Definable compactness and definable subgroups of o-minimal groups*, J. London Math. Soc. **59** (1999), 769–786.
- [9] M. Shiota, *Geometry of subanalytic and semialgebraic sets*, Progress in Math. **150** (1997), Birkhäuser.

[1] E. Baro and M. Otero, *On o-minimal*