

# An affine definable $C^rG$ manifold admits a unique affine definable $C^\infty G$ manifold structure

Tomohiro Kawakami

Department of Mathematics, Faculty of Education, Wakayama University,  
Sakaedani, Wakayama 640-8510, Japan  
kawa@center.wakayama-u.ac.jp

Received July 7, 2017

## Abstract

Let  $G$  be a compact subgroup of  $GL_n(\mathbb{R})$ . We prove that every affine definable  $C^rG$  manifold admits a unique affine definable  $C^\infty G$  manifold structure up to definable  $C^\infty G$  diffeomorphism ( $1 \leq r < \infty$ ). Moreover we prove that every strongly definable  $C^rG$  vector bundle over  $X$  admits a unique strongly definable  $C^\infty G$  vector bundle structure up to definable  $C^\infty G$  vector bundle isomorphism ( $0 \leq r < \infty$ ). Furthermore we consider raising differentiability of strong definable  $C^r$  fiber bundles ( $0 \leq r < \infty$ ).

2010 *Mathematics Subject Classification.* 57S15, 14P20, 57R35, 58A07, 03C64.

*Keywords and Phrases.* Definable  $C^\infty G$  manifolds, definable  $C^\infty G$  maps, approximation theorem, definable  $C^\infty G$  vector bundles, definable  $C^\infty$  fiber bundles, o-minimal.

## 1. Introduction.

By [15], if  $s$  is a non-negative integer, then every  $C^s$  Nash map between affine Nash manifolds is approximated in the definable  $C^s$  topology by Nash maps. This definable  $C^s$  topology is a new topology defined in [15]. There is a generalization of this result in the definable  $C^r$  category obtained by an o-minimal expansion  $\mathcal{M} = (\mathbb{R}, +, \cdot, <, \dots)$  on the standard structure  $\mathcal{R} = (\mathbb{R}, +, \cdot, <)$  of the field  $\mathbb{R}$  of real numbers, namely if  $0 \leq s < r < \infty$ , then every definable  $C^s$  map between affine definable  $C^r$  manifolds is approximated in the definable  $C^s$  topology by definable  $C^r$  maps (II.5.2 [16]).

In this paper,  $G$  denotes a compact subgroup of  $GL_n(\mathbb{R})$ , every definable map is con-

tinuous and any manifold does not have boundary, unless otherwise stated. Under our assumption,  $G$  is a compact algebraic subgroup of  $GL_n(\mathbb{R})$  (e.g. 2.2 [13]). We consider an equivariant definable version of the above theorem in an o-minimal expansion  $\mathcal{M}$  and an affine definable  $C^\infty G$  manifold structure of an affine definable  $C^rG$  manifold. General references on o-minimal structures are [2], [4], see also [16]. Further properties and constructions of them are studied in [3], [5], [14].

We also consider strongly definable  $C^\infty G$  vector bundle structures of strongly definable  $C^rG$  vector bundles ( $0 \leq r < \infty$ ). Moreover we consider raising differentiability of strong definable  $C^r$  fiber bundles ( $0 \leq r < \infty$ ).

Suppose that  $\eta$  is a definable  $C^r G$  vector bundle over an affine definable  $C^r G$  manifold  $X$  and  $0 \leq r \leq \infty$ . We say that  $\eta$  is *strongly definable* if there exist a representation  $\Omega$  of  $G$  and a definable  $C^r G$  map  $f : X \rightarrow G(\Omega, \alpha)$  such that  $\eta$  is definably  $C^r G$  vector bundle isomorphic to  $f^*(\gamma(\Omega, \alpha))$ , where  $\alpha$  denotes the rank of  $\eta$ .

The following is the main result of this paper.

**Theorem 1.1.** *Let  $X$  be an affine definable  $C^r G$  manifold and  $\mathcal{M}$  admits  $C^\infty$  cell decomposition and exponential.*

(1) *If  $1 \leq r < \infty$  then,  $X$  admits a unique affine definable  $C^\infty G$  manifold structure up to definable  $C^\infty G$  diffeomorphism.*

(2) *If  $0 \leq r < \infty$ , then every strongly definable  $C^r G$  vector bundle over an affine definable  $C^\infty G$  manifold admits a unique strongly definable  $C^\infty G$  vector bundle structure up to definable  $C^\infty G$  vector bundle isomorphism.*

(3) *If  $0 \leq r < \infty$ , each strongly definable  $C^r$  fiber bundle over an affine definable  $C^\infty$  manifold admits a unique strongly definable  $C^\infty$  fiber bundle structure up to definable  $C^\infty$  fiber bundle isomorphism.*

Remark.

(1) If  $1 \leq s < r < \infty$ , then definable  $C^r$  manifold structures of definable  $C^s$  manifolds are studied in [8].

(2) If  $1 \leq s < r < \infty$  and  $G$  is finite, then strongly definable  $C^r$  vector bundle structures of strongly definable  $C^s G$  vector bundles are studied in [9].

(3) If  $1 \leq s < r < \infty$ , then strongly definable  $C^r$  fiber bundle structures of strongly definable  $C^s$  fiber bundles are studied in [6].

## 2. Definable $C^r G$ manifolds.

Recall the definition of definable  $C^r G$  manifolds ([11], [9]).

**Definition 2.1** ([11], [9]). Let  $0 \leq r \leq \infty$ .

- (1) A group homomorphism (resp. A group isomorphism) from  $G$  to  $O_n(\mathbb{R})$  is a *definable group homomorphism* (resp. a *definable group isomorphism*) if it is a definable map (resp. a definable homeomorphism).

Note that a definable group homomorphism (resp. a definable group isomorphism) between  $G$  and  $O_n(\mathbb{R})$  is a definable  $C^\infty$  map (resp. a definable  $C^\infty$  diffeomorphism) because  $G$  and  $O_n(\mathbb{R})$  are Lie groups.

- (2) An *n-dimensional representation* of  $G$  means  $\mathbb{R}^n$  with the linear action induced by a definable group homomorphism from  $G$  to  $O_n(\mathbb{R})$ . In this paper, we assume that every representation of  $G$  is orthogonal.
- (3) A *definable  $C^r G$  manifold* is a pair  $(X, \alpha)$  consisting of a definable  $C^r$  manifold  $X$  and a group action  $\alpha$  of  $G$  on  $X$  such that  $\alpha : G \times X \rightarrow X$  is a definable  $C^r$  map. For simplicity of notation, we write  $X$  instead of  $(X, \alpha)$ .
- (4) A definable  $C^r$  submanifold of a definable  $C^r G$  manifold  $X$  is called a *definable  $C^r G$  submanifold* of  $X$  if it is  $G$  invariant.
- (5) A definable  $C^r$  map (resp. A definable  $C^r$  diffeomorphism, A definable homeomorphism, A definable map) is a *definable  $C^r G$  map* (resp. a *definable  $C^r G$  diffeomorphism*, a *definable  $G$  homeomorphism*, a *definable  $G$  map*) if it is a  $G$  map.
- (6) A definable  $C^r G$  manifold is called *affine* if it is definably  $C^r G$  diffeomorphic (definably  $G$  homeomorphic if  $r = 0$ ) to a definable  $C^r G$  submanifold of some representation of  $G$ .
- (7) A *definable  $C^r G$  manifold with boundary* is defined similarly.

If  $0 \leq r < \infty$ , then every definable  $C^r$  manifold is affine ([11], [10]) and if  $\mathcal{M}$  is exponential, then each compact definable  $C^\infty G$  manifold is affine [11].

Recall the definable  $C^s$  topology [9] and some results on it [9].

Let  $X$  and  $Y$  be definable  $C^s$  submanifolds of  $\mathbb{R}^n$  and  $\mathbb{R}^m$ , respectively, and  $0 \leq s < \infty$ . Let  $C_{def}^s(X, Y)$  denote the set of definable  $C^s$  maps from  $X$  to  $Y$ . For  $f \in C_{def}^s(X, Y)$  and  $x \in X$ , the differential  $df_x$  of  $f$  at  $x$  means a linear map from the tangent space  $T_x X$  of  $X$  at  $x$  to  $\mathbb{R}^m$ . Composing it with the orthogonal projection  $\mathbb{R}^n \rightarrow T_x X$ , one can extend  $df_x$  to a linear map  $\mathbb{R}^n \rightarrow \mathbb{R}^m$ . Then  $Df : X \rightarrow M(m, n; \mathbb{R}) = \mathbb{R}^{mn}$  is defined as the matrix representation of  $df$ . For each  $1 \leq k \leq s$ , we inductively define a  $C^{s-k}$  map

$$D^k f : X \rightarrow \mathbb{R}^{n^k m}, D^k f = D(D^{k-1} f).$$

Let  $\|f\|_s$  denote the definable function on  $X$  defined by

$$\|f\|_s(x) = |f(x)| + |Df(x)| + \cdots + |D^s f(x)|.$$

For a positive definable function  $\epsilon : X \rightarrow \mathbb{R}$ , let

$$U_\epsilon = \{h \in C_{def}^s(X, Y) \mid \|h\|_s < \epsilon\}.$$

We say that the *definable  $C^s$  topology* on  $C_{def}^s(X, Y)$  is the topology defined by choosing  $\{h + U_\epsilon\}_\epsilon$  as a fundamental neighborhood system of  $h$  in  $C_{def}^s(X, Y)$ . In the Nash category, we simply call it the  *$C^r$  topology*. If  $X$  is compact, then this topology coincides with the  $C^s$  Whitney topology (p 156 [16]).

**Proposition 2.2** ([16], 4.9 [9]). *Let  $X$ ,  $Y$  and  $Z$  be definable  $C^s$  submanifolds  $\mathbb{R}^n$ ,  $\mathbb{R}^m$  and  $\mathbb{R}^l$ , respectively, and  $0 \leq s < \infty$ . Let  $f \in C_{def}^s(X, Y)$  and  $h \in C_{def}^s(Y, Z)$ .*

- (1) *The map  $h_* : C_{def}^s(X, Y) \rightarrow C_{def}^s(X, Z)$ ,  $h_*(k) = h \circ k$  is continuous.*
- (2) *The map  $f^* : C_{def}^s(Y, Z) \rightarrow C_{def}^s(X, Z)$ ,  $f^*(k) = k \circ f$  is continuous if and only if  $f$  is proper.*

**Proposition 2.3** ([16], 4.10 [9]). *Let  $X$  and  $Y$  be definable  $C^s$  submanifolds of  $\mathbb{R}^n$  and  $0 < s < \infty$ . Let  $f : X \rightarrow Y$  be a definable  $C^s$  map. If  $f$  is an immersion (resp. a*

*diffeomorphism, a diffeomorphism onto its image), then an approximation of  $f$  in the definable  $C^s$  topology is an immersion (resp. a diffeomorphism, a diffeomorphism onto its image). Moreover if  $f$  is a diffeomorphism, then  $h^{-1} \rightarrow f^{-1}$  as  $h \rightarrow f$ .*

**Theorem 2.4** ([16], 4.11 [9]). *Let  $X$  and  $Y$  be affine definable  $C^r$  manifolds and  $0 \leq s < r < \infty$ . Then every definable  $C^s$  map  $f : X \rightarrow Y$  is approximated in the definable  $C^s$  topology by definable  $C^r$  maps.*

**Theorem 2.5** ([8]). *If  $0 \leq s < r < \infty$ , then every definable  $C^s G$  map between affine definable  $C^r G$  manifolds is approximated in the definable  $C^s$  topology by definable  $C^r G$  maps.*

**Proposition 2.6** ([11]). *Every definable  $C^\infty G$  submanifold  $X$  of a representation  $\Omega$  of  $G$  has a definable  $C^\infty G$  tubular neighborhood  $(U, \theta)$  of  $X$  in  $\Omega$ , namely  $U$  is a  $G$  invariant definable open neighborhood of  $X$  in  $\Omega$  and  $\theta : U \rightarrow X$  is a definable  $C^\infty G$  map with  $\theta|_X = id_X$ .*

**Proposition 2.7** ([8]). *(Equivariant definable  $C^r$  partition of unity). Let  $X$  be a definable  $C^r G$  submanifold closed in a representation  $\Omega$  of  $G$  and  $\{U_i\}_{i=1}^l$  a finite  $G$  invariant definable open covering of  $X$  and  $0 \leq r < \infty$ . Then there exist  $G$  invariant definable  $C^r$  functions  $\lambda_1, \dots, \lambda_l : X \rightarrow \mathbb{R}$  such that  $0 \leq \lambda_i \leq 1$ ,  $\text{supp } \lambda_i \subset U_i$  and  $\sum_{i=1}^l \lambda_i(x) = 1$  for any  $x \in X$ .*

By a way similar to the proof of the above proposition, we have the following.

**Proposition 2.8** (Equivariant definable  $C^\infty$  partition of unity). *Suppose that  $\mathcal{M}$  admits  $C^\infty$  cell decomposition and exponential. Let  $X$  be a definable  $C^\infty G$  submanifold closed in a representation  $\Omega$  of  $G$  and  $\{U_i\}_{i=1}^l$  a finite  $G$  invariant definable open covering of  $X$ . Then there exist  $G$  invariant definable  $C^\infty$  functions  $\lambda_1, \dots, \lambda_l : X \rightarrow \mathbb{R}$  such that  $0 \leq \lambda_i \leq 1$ ,  $\text{supp } \lambda_i \subset U_i$  and  $\sum_{i=1}^l \lambda_i(x) = 1$  for any  $x \in X$ .*

**Proposition 2.9** ([9]). *Let  $X$  be a compact affine definable  $C^\infty G$  manifold with boundary  $\partial X$ . Then  $X$  admits a definable  $C^\infty G$  collar, that is, there exists a definable  $C^\infty G$  imbedding  $\phi : \partial X \times [0, 1) \rightarrow X$  such that  $\phi|_{\partial X \times \{0\}} = id_{\partial X}$ , where the action on  $[0, 1)$  is trivial.*

**Theorem 2.10** ([9]). *Every definable  $C^\infty G$  manifold is either compact or compactifiable.*

Let  $f$  be a map from a  $C^r G$  manifold  $X$  to a representation  $\Omega$  of  $G$  and  $0 \leq r \leq \infty$ . Denote the Haar measure of  $G$  by  $dg$ , and let  $x$  be a point in  $X$ . Recall the averaging operator  $A$  is defined by

$$A(f)(x) = \int_G g^{-1} f(gx) dg.$$

**Proposition 2.11** (4.1 [1]). *Let  $G$  be a compact Lie group and  $0 \leq r \leq \infty$ . Suppose that  $C^r(X, \Omega)$  denotes the set of  $C^r$  maps from a  $C^r G$  submanifold  $X$  of a representation of  $G$  to a representation  $\Omega$  of  $G$ .*

- (1) *The averaged map  $A(f)$  of  $f$  is equivariant, and  $A(f) = f$  if  $f$  is equivariant.*
- (2) *If  $f \in C^r(X, \Omega)$ , then  $A(f) \in C^r(X, \Omega)$ .*
- (3) *If  $f$  is a polynomial map, then so is  $A(f)$ .*
- (4) *If  $X$  is compact and  $r < \infty$ , then  $A : C^r(X, \Omega) \rightarrow C^r(X, \Omega)$  is continuous in the  $C^r$  Whitney topology.*

**Theorem 2.12.** *If  $0 \leq s < \infty$  and  $M$  admits  $C^\infty$  cell decomposition and exponential, then every definable  $C^s G$  map between affine definable  $C^\infty G$  manifolds is approximated in the definable  $C^s$  topology by definable  $C^\infty G$  maps.*

*Proof.* Let  $f : X \rightarrow Y$  be a definable  $C^s G$  map. If  $X$  is compact, the proof is easy. We assume that  $X$  is noncompact.

Since  $M$  admits  $C^\infty$  cell decomposition, is exponential and by Theorem 2.10,  $X$  is definably  $C^\infty G$  diffeomorphic to the interior of

a compact definable  $C^\infty G$  manifold  $Y$  with boundary  $\partial Y$ . By Proposition 2.9, we can take the double  $W$  of  $Y$ . Then  $W$  is a compact definable  $C^\infty G$  manifold. By [11],  $W$  is affine. Note that  $X$  is a definable  $C^\infty G$  submanifold of  $W$ .

Since  $M$  admits  $C^\infty$  cell decomposition and by 2.3 [9], there exists a definable open subset  $Z$  of  $X$  such that  $\dim(X - Z) < \dim X$  and  $f|_Z$  is a definable  $C^\infty$  map. Since  $X$  is a definable  $G$  set, we can take  $Z$  which is definable and  $G$  invariant. Since the action is orthogonal, the  $\epsilon$  neighborhood  $N(Z, \epsilon) = \{x \in X | d(x, Z) < \epsilon\}$  is a  $G$  invariant definable open  $G$  set. Since  $W$  is compact,  $N(Z, \epsilon)$  is bounded and the closure  $N'$  of  $N(Z, \epsilon)$  is compact. Applying Proposition 2.11, there exists a polynomial  $G$  map  $F : N' \rightarrow \Xi$  such that  $F$  is an approximation of  $f|_{N'}$ . By Proposition 2.8, gluing  $f|_{X - Z}$  and  $f|_{N'}$ , we have a definable  $C^\infty G$  map  $h : X \rightarrow \Xi$ . By Proposition 2.6, there exists a definable  $C^\infty G$  tubular neighborhood  $(U, \theta)$  of  $Y$  in  $\Xi$ . The map defined by  $\theta \circ h$  is the required map.  $\square$

By a way to a partial proof of equivariant Nash conjecture, we have the following theorem.

**Theorem 2.13.** *Let  $X$  be an affine definable  $C^r G$  manifold and  $1 \leq r < \infty$ . Then  $X$  admits an affine definable  $C^\infty G$  manifold structure.*

*Proof of Theorem 1.1 (1).* By Theorem 2.13,  $X$  admits an affine definable  $C^\infty G$  manifold structure. Uniqueness of affine definable  $C^\infty G$  manifold structure follows from Theorem 2.12 and Proposition 2.3.  $\square$

Remark that nonaffine definable  $C^\infty G$  manifold structures of an affine definable  $C^r G$  manifold is not necessarily unique even if  $M = (\mathbb{R}, +, \cdot, <)$  ([12]). If the action on  $X$  is transitive, then definable  $C^\infty G$  manifold structure is unique and there is no nonaffine definable  $C^\infty G$  manifold structure ([12]).

### 3. Definable $C^rG$ vector bundles.

Recall the definition of definable  $C^rG$  vector bundles [9].

**Definition 3.1** ([9]). Suppose that  $0 \leq r \leq \infty$ .

- (1) A *definable  $C^rG$  vector bundle* is a definable  $C^r$  vector bundle  $\eta = (E, p, X)$  satisfying the following three conditions.
  - (a) The total space  $E$  and the base space  $X$  are definable  $C^rG$  manifolds.
  - (b) The projection  $p : E \rightarrow X$  is a definable  $C^rG$  map.
  - (c) For any  $x \in X$  and  $g \in G$ , the map  $p^{-1}(x) \rightarrow p^{-1}(gx)$  is linear.
- (2) Let  $\eta$  and  $\zeta$  be definable  $C^rG$  vector bundles over  $X$ . A definable  $C^r$  vector bundle morphism  $\eta \rightarrow \zeta$  is called a *definable  $C^rG$  vector bundle morphism* if it is a  $G$  map. A definable  $C^rG$  vector bundle morphism  $f : \eta \rightarrow \zeta$  is said to be a *definable  $C^rG$  vector bundle isomorphism* if there exists a definable  $C^rG$  vector bundle morphism  $h : \zeta \rightarrow \eta$  such that  $f \circ h = id$  and  $h \circ f = id$ .
- (3) A definable  $C^r$  section of a definable  $C^rG$  vector bundle is a *definable  $C^rG$  section* if it is a  $G$  map.
- (4) If  $r = 0$ , then a definable  $C^0G$  vector bundle (resp. a definable  $C^0G$  vector bundle morphism, a definable  $C^0G$  vector bundle isomorphism, a definable  $C^0G$  section) is simply called a *definable  $G$  vector bundle* (resp. a *definable  $G$  vector bundle morphism*, a *definable  $G$  vector bundle isomorphism*, a *definable  $G$  section*).

Recall universal  $G$  vector bundles (e.g. [9]) and existence of a Nash  $G$  tubular neighborhood of a Nash  $G$  submanifold of a representation of  $G$  (2.3 [12]).

**Definition 3.2.** Let  $\Omega$  be an  $n$ -dimensional representation of  $G$  induced by a definable group homomorphism  $B : G \rightarrow O_n(\mathbb{R})$ . Suppose that  $M(\Omega)$  denotes the vector space of  $n \times n$ -matrices with the action  $(g, A) \in G \times M(\Omega) \rightarrow B(g)AB(g)^{-1} \in M(\Omega)$ . For any positive integer  $\alpha$ , we define the vector bundle  $\gamma(\Omega, \alpha) = (E(\Omega, \alpha), u, G(\Omega, \alpha))$  as follows:  $G(\Omega, \alpha) = \{A \in M(\Omega) | A^2 = A, A = A', TrA = \alpha\}$ ,  $E(\Omega, \alpha) = \{(A, v) \in G(\Omega, \alpha) \times \Omega | Av = v\}$ ,  $u : E(\Omega, \alpha) \rightarrow G(\Omega, \alpha)$ ,  $u((A, v)) = A$ , where  $A'$  denotes the transposed matrix of  $A$  and  $Tr A$  stands for the trace of  $A$ . Then  $\gamma(\Omega, \alpha)$  is an algebraic vector bundle. Since the action on  $\gamma(\Omega, \alpha)$  is algebraic, it is an algebraic  $G$  vector bundle. We call it *the universal  $G$  vector bundle associated with  $\Omega$  and  $\alpha$* . Remark that  $G(\Omega, \alpha) \subset M(\Omega)$  and  $E(\Omega, \alpha) \subset M(\Omega) \times \Omega$  are nonsingular algebraic  $G$  sets. In particular, they are Nash  $G$  submanifolds of  $M(\Omega)$  and  $M(\Omega) \times \Omega$ , respectively.

**Definition 3.3** ([9]). (1) Let  $G$  be a definable group. A definable  $G$  vector bundle  $\eta = (E, p, X)$  over a definable  $G$  set  $X$  is called *strongly definable* if there exist a representation  $\Omega$  of  $G$  and a definable  $G$  map  $f : X \rightarrow G(\Omega, k)$  such that  $\eta$  is definably  $G$  vector bundle isomorphic to  $f^*(\gamma(\Omega, k))$ , where  $k$  denotes the rank of  $\eta$ .

(2) Let  $G$  be a definable  $C^r$  group and  $0 \leq r \leq \infty$ . A definable  $C^rG$  vector bundle  $\eta = (E, p, X)$  over an affine definable  $C^rG$  manifold  $X$  is called *strongly definable* if there exist a representation  $\Omega$  of  $G$  and a definable  $C^rG$  map  $f : X \rightarrow G(\Omega, k)$  such that  $\eta$  is definably  $C^rG$  vector bundle isomorphic to  $f^*(\gamma(\Omega, k))$ , where  $k$  denotes the rank of  $\eta$ .

**Proposition 3.4** (2.3 [12]). *Every Nash  $G$  submanifold  $X$  of a representation  $\Omega$  of  $G$  has a Nash  $G$  tubular neighborhood  $(U, \theta)$  of  $X$  in  $\Omega$ .*

*Proof of Theorem 1.1* (2) Let  $\eta$  be a strongly definable  $C^rG$  vector bundle over  $X$ . Since  $\eta$  is strongly definable, there exists a definable  $C^rG$  map  $f : X \rightarrow G(\Omega, \alpha)$

such that  $\eta$  is definably  $C^rG$  vector bundle isomorphic to  $f^*(\gamma(\Omega, \alpha))$ .

By Theorem 1.1 (1),  $f$  is approximated by a definable  $C^\infty G$  map  $h : X \rightarrow G(\Omega, \alpha)$ . By [9],  $\eta$  is definably  $C^rG$  vector bundle isomorphic to a strongly definable  $C^\infty G$  vector bundle  $h^*(\gamma(\Omega, \alpha))$ . By 1.7 [9] and Theorem 1.1 (1), uniqueness follows.  $\square$

## 4. Definable $C^r$ fiber bundles.

By a way similar to the proof of 2.6 [6], we have the following.

**Proposition 4.1.** *Let  $\mathcal{B}_K = (B_K, p_K, X_K)$  be the  $n$ -universal principal bundle relative to  $K$ ,  $F$  an affine definable  $C^\infty$  manifold with an effective definable  $C^\infty K$  action. Then the associated fiber bundle  $\mathcal{B}_K[F] := (E, p, X_K, F, K)$  is a definable  $C^\infty$  fiber bundle.*

*Proof of Theorem 1.1 (3).* Let  $\eta$  be a strongly definable  $C^r$  fiber bundle over  $X$ . Then there exists the  $n$ -universal bundle  $\mathcal{B}_K$  and a definable map  $f : X \rightarrow X_K$  such that  $f^*(\mathcal{B}_K[F])$  is definably fiber bundle isomorphic to  $\eta$ . By Theorem 2.12, we have a definable  $C^\infty$  map  $h : X \rightarrow X_K$  as an approximation of  $f$ . In particular  $h$  is definably homotopic to  $f$ . Thus by 1.1 [7],  $\zeta := h^*(\mathcal{B}_K[F])$  is definably fiber bundle isomorphic to  $f^*(\mathcal{B}_K[F])$  and  $\zeta$  is a strongly definable  $C^\infty$  fiber bundle.

(2) Let  $\zeta'$  be another strongly definable  $C^\infty$  fiber bundle over  $X$  such that  $\zeta'$  is definably fiber bundle isomorphic to  $\eta$ . Consider the strongly definable  $C^r$  fiber bundle  $(\zeta, \zeta', id_X)$  whose sections represent the fiber bundle isomorphisms between  $\zeta$  and  $\zeta'$  which is defined in 2.11 [6]. Then it has a continuous section. By a way similar to the proof of 2.12 [6], it admits a definable  $C^\infty$  section. This section gives a definable  $C^\infty$  fiber bundle isomorphism between  $\zeta$  and  $\zeta'$ .  $\square$

## References

- [1] K.H. Dovermann, M. Masuda, and T. Petrie, *Fixed point free algebraic actions on varieties diffeomorphic to  $\mathbb{R}^n$* , Progress in Math. **80**, Birkhäuser (1990), 49–80.
- [2] L. van den Dries, *Tame topology and o-minimal structures*, Lecture notes series **248**, London Math. Soc. Cambridge Univ. Press (1998).
- [3] L. van den Dries, A. Macintyre, and D. Marker, *The elementary theory of restricted analytic field with exponentiation*, Ann. of Math. **140** (1994), 183–205.
- [4] L. van den Dries and C. Miller, *Geometric categories and o-minimal structures*, Duke Math. J. **84** (1996), 497–540.
- [5] L. van den Dries and P. Speissegger, *The real field with convergent generalized power series*, Trans. Amer. Math. Soc. **350**, (1998), 4377–4421.
- [6] T. Kawakami, *Definable  $C^r$  fiber bundles and definable  $C^rG$  vector bundles*, Commun. Korean Math. Soc. **23** (2008), 257–268.
- [7] T. Kawakami, *Definable fiber bundles in an o-minimal expansion of a real closed field*, Bull. Fac. Ed. Wakayama Univ. Natur. Sci. No. **60** (2010), 15–23.
- [8] T. Kawakami, *Equivariant definable  $C^r$  approximation theorem, definable  $C^rG$  triviality of  $G$  invariant definable  $C^r$  functions and compactifications*, Bull. Fac. Ed. Wakayama Univ. Natur. Sci. **55** (2005), 23–36.
- [9] T. Kawakami, *Equivariant differential topology in an o-minimal expansion of the field of real numbers*, Topology and its appl. **123** (2002), 323–349.
- [10] T. Kawakami, *Every definable  $C^r$  manifold is affine*, Bull. Korean Math. Soc. **42** (2005), 165–167.

- [11] T. Kawakami, *Imbedding of manifolds defined on an o-minimal structures on  $(\mathbb{R}, +, \cdot, <)$* , Bull. Korean Math. Soc. **36** (1999), 183–201.
- [12] T. Kawakami, *Nash  $G$  manifold structures of compact or compactifiable  $C^\infty G$  manifolds*, J. Math. Soc. Japan **48** (1996), 321–331.
- [13] D.H. Park and D.Y. Suh, *Linear embeddings of semialgebraic  $G$ -spaces*, Math. Z. **242**, (2002), 725–742.
- [14] Y. Peterzil, A. Pillay and S. Starchenko, *Definably simple groups in o-minimal structures*, Trans. Amer. Math. Soc. **352** (2000), 4397–4419.
- [15] M. Shiota, *Approximation theorems for Nash mappings and Nash manifolds*, Trans. Amer. Math. Soc. **293** (1986), 319–337.
- [16] M. Shiota, *Geometry of subanalytic and semialgebraic sets*, Progress in Math. **150** (1997), Birkhäuser.

